



CFG  
Law

part of the client first group

## **Important information about sustaining a concussion and mild Traumatic Brain Injury**



# Contents

Introduction	3
About mild TBIs	4 - 5
Vital signs to look out for after a head injury	6
What are the symptoms of a mild TBI?	7
Cognitive symptoms	8 - 11
Physical symptoms	12 - 16
Emotional symptoms	17 - 19
Post-Concussion Syndrome	20
Diagnosing mild brain injuries	21
How a mild brain injury can impact your life	22 - 25
Further support after a mild brain injury	26
Charities and support groups	27
Helping friends and family to understand brain injuries	28
About CFG Law	29



# Introduction

Brain injuries are often referred to as ‘invisible injuries’, because it’s impossible for others to fully understand what’s happening below the surface. While there are a lot of common symptoms and issues associated with brain injuries, every case is unique.

Because each person’s journey to recovery takes a different path, after a mild brain injury (such as concussion), it can be hard to see the big picture of the obstacles you may face, and the treatment and support options available.

This simple guide is designed to offer you useful, easy to understand information about mild Traumatic Brain Injury (TBI) - from the immediate impact of the injury to the potential affect it can have on your life.



# About mild TBIs



Mild Traumatic Brain Injuries (TBIs) are usually caused by a knock, blow or jolt to the head that results in a temporary disturbance to the brain's functionality.

Concussion is the most common type of mild TBI, with symptoms usually lasting only a few days or weeks, although some people can experience long term symptoms.

Medical professionals class a brain injury as 'mild' if a person experiences a loss of consciousness or disorientation for less than 30 minutes.

Mild TBIs usually don't show up on CT or MRI scans, making them difficult to readily diagnose and understand. Most people are discharged from hospital quickly and will experience no long-term damage to the brain. It's not uncommon, however, for 'mild' injuries to be more serious than first thought.

Often the seriousness of ‘mild’ TBIs isn’t obvious to either medical staff or the injured person themselves, although family members are often the first to recognise something is not right. While some people recover from a mild TBI quickly, others can experience lasting effects from their injury.

[Researchers have found that about 15% of patients](#) who are diagnosed with a mild TBI continue to suffer serious symptoms long after their initial injury. Unfortunately, many people underestimate or are unaware of the full extent of their TBI, and try to get back to normal life too quickly, resulting in more serious problems.

**That’s why it’s important to take your symptoms seriously, and be guided by medical professionals on your road to recovery.**



# Vital signs to look out for after a head injury

If you are experiencing any of the following symptoms in the days after sustaining a head injury, you should dial 111 or go to your nearest A&E department as you may have sustained a brain injury:

Vomiting	Convulsions or fits	Collapsing or suddenly passing out	Unconsciousness or trouble staying conscious (e.g. problems keeping your eyes open)
----------	---------------------	------------------------------------	---

Any confusion (e.g. not knowing where you are and getting things mixed up)	Problems with speech or understanding conversations	Clear fluid or blood coming out of ears or nose	New deafness in one or both ears	Problems with balance or walking
--	---	---	----------------------------------	----------------------------------

Weakness in arms or legs	Problems with sight	Severe headache that is not relieved by painkillers	Unusual drowsiness when you'd normally be wide awake
--------------------------	---------------------	---	--

# What are the symptoms of a mild TBI?

Immediately after suffering a brain injury, it's likely you'll experience some of the following symptoms:



Mild Headache



Feeling nauseas  
(without being sick)



Dizziness



Irritability and  
bad temper



Concentration issues  
and problems with  
memory



Tiredness



Lack of  
appetite



Sleep problems  
such as disturbed  
sleep

These are some of the more common symptoms people with brain injuries experience, but there are a wide range of different issues associated with mild TBIs too. These are usually divided into three categories: cognitive, physical and emotional symptoms. Let's have a look at each category.

# Cognitive symptoms

‘Cognitive’ refers to the way we think and process information. A brain injury can affect how a person thinks, learns and remembers. Often, when somebody experiences a brain injury, they can get stressed or worried about the changes they’re experiencing. While this is completely understandable, it can make these problems worse.

Here are some of the most common cognitive symptoms you may be struggling with:

Memory  
problems

Poor  
concentration  
and reduced  
attention span

Behaving  
inappropriately  
and oversharing

# Memory problems

Poor short term memory is one of the most common issues associated with brain injuries. Struggling to remember things can be frustrating, both for you and your loved ones.



There are also a few things you can do to make life a little easier for you and your family as you recover. Using post-it notes, calendars and diaries can help remind you to do important things, such as lock the door, remember your wallet when you leave the house, or attend appointments.

It can also be useful to carry a notebook and pen with you to write down anything you need to remember. Tidying things away to a specific place is also a good tip for remembering where things are stored around the house.

# Concentration and attention span



It's also common for people to struggle with concentration and attention span after a brain injury. This makes it particularly difficult to focus in loud environments, or concentrate on two things at once. A brain injury may impact your concentration span, making it hard to stay alert at work or school, which often affects your performance.

If you're struggling with concentration, try to do any important tasks when you know you are likely to be at your most alert.

It's also a good idea to reduce background noise where possible, and complete anything you want to get done somewhere quiet. Try doing only one task at a time, and slowly build your concentration levels back up to avoid getting frustrated.



# Behaving inappropriately and oversharing

Brain injury can also impact the way people behave. Common issues include a loss of inhibitions (called disinhibition), leading to inappropriate behaviour. After a mild TBI, you may make unacceptable remarks or laugh inappropriately, which can upset your loved ones.

Oversharing of personal information can also be common, and again can upset people around you or make them feel uncomfortable.

Most people will gradually regain control of their behaviour, but sometimes these disinhibitions will remain long term. You should try and consider the comments made by loved ones about your behaviour and think about how it may be perceived by other people.



# Physical symptoms

As well as problems with thinking and processing information, you may also be struggling with a number of physical issues too. Because the brain is so central to how we function, any injuries to your head can cause problems with everyday tasks.

Like the symptoms above, physical problems are likely to get better over time - but it's impossible to know how quickly you will recover. Try to stay patient. It is a good idea to ease yourself back into physical activities, and listen to your body!

Sleep  
problems

Fatigue

Headaches

Dizziness and  
balance  
problems

# Sleep problems

After a brain injury, you may experience problems with sleeping. While it can be difficult, it's important that you get enough rest after a brain injury to help your recovery. In the first few weeks, this might be much more than you'd usually need, and even after this initial period has passed, it's still important to get lots of sleep.

Try sticking to a regular sleep routine by going to bed and waking up at the same time every day. Even if you're sleeping well at night, many people need naps to avoid exhaustion. If you do nap, set alarms so you're not snoozing for too long - as this can have an impact on the quality of sleep you get at night.

Stress and worry can also affect sleep. If you're struggling with this, then it's a good idea to try relaxation techniques such as meditation.



# Fatigue



Fatigue is not necessarily linked to sleep, as brain injuries can cause extreme tiredness even in people who are sleeping well at night. Feeling exhausted can have a big impact on your daily life, and make normal tasks difficult.

If you keep a diary about your experiences of fatigue, it might be easier to identify any common triggers. If you know what is causing tiredness, you can either avoid these activities if possible, or learn to manage them better in the future.

It's also a good idea to prioritise important tasks when you have more energy. Break your daily routine down into manageable chunks and leave plenty of time to rest in between.

Although it can be frustrating, try not to push yourself too much and recognise when you need to rest. If you try and push through fatigue, it can cause more problems and make the recovery time longer. It's important to ease yourself back into your daily tasks at a pace that suits you.

# Headaches

Headaches are a common symptom of mild brain injuries like concussions, particularly in the immediate aftermath of the injury. If your headache is still there after several days and isn't relieved by painkillers, you should seek medical attention to check for more serious complications.

As you continue to recover, headaches can be a reaction to other symptoms - such as fatigue, stress and worry, or concentration and memory issues. As with fatigue, you should try to keep a diary of your headaches to identify what triggers them.

Although headaches can often be very uncomfortable, try not to become too reliant on painkillers - as long-term use of certain medications can be addictive, which actually makes headaches worse.



# Dizziness and balance problems

Brain injuries can cause problems with balance and lead to feelings of dizziness. You may find that you're unsteady on your feet and have problems with coordination, often bumping into things.

These symptoms will usually pass quite quickly, but make sure to take extra care early on, particularly when crossing the road. You should never drive if you have any balance problems or are experiencing dizziness.

You should also try to keep things tidy around your home and remove any obvious tripping hazards and obstacles until this passes. It may be necessary to introduce grab rails on stairs and in bathrooms, or use a perching stools to complete household tasks to avoid prolonged standing.



# Emotional symptoms

After suffering a mild brain injury, it's normal for your emotions to be all over the place. Feelings of frustration, sadness, confusion and anxiety are all common and completely understandable. It's often difficult to express your feelings after a brain injury, and for your family and friends to fully understand what you're going through.

The important thing to remember is that your loved ones almost certainly mean well and are trying to help you get better - even if you find yourself clashing with them. If you want to talk to somebody, you can find information on charities who can help you at the end of this guide.

Anger and  
irritability

Low mood,  
depression and  
anxiety

# Anger and irritability

Feelings of anger, losing your temper and irritability are common after a brain injury. You can often find yourself getting agitated and annoyed by things that previously wouldn't have bothered you. This may be partly because of other symptoms you are experiencing, such as frustration at memory problems, or being annoyed with ongoing fatigue and headaches.



To reduce irritability, rest as much as you can, follow a routine that's familiar to you, and try to recognise anything that triggers your anger. As we mentioned earlier, other things like meditation and relaxation techniques can also help you to control your emotions. If you feel yourself getting angry, try to take some time out to calm down.

# Low mood, depression and anxiety

Sustaining a brain injury can be stressful and worrying. It's difficult trying to come to terms with the various symptoms you may be experiencing, which can lead to low mood, as well as feelings of depression and anxiety.



The physical and cognitive symptoms we've looked at can take their toll, and it's normal to feel emotional after a mild brain injury. You can speak to your GP about how you're feeling, and they may be able to refer you to more specialist treatment from a neuropsychologist or healthcare professional who specialises in depression after a brain injury.

After a head injury, a lot of these emotions are a result of trauma to the brain, and may be completely irrational. If you do find that you're a lot more emotional, you shouldn't feel guilty about any outbursts! Understanding the reasons for your emotions will help you to explain them to your family, so they don't take it personally.

# Post-Concussion Syndrome

If any other ongoing symptoms are still there after a long period of time, you may be diagnosed with Post-Concussion Syndrome (PCS). Normally, symptoms of a mild brain injury should subside within two weeks. If you are still struggling after this time, you should seek advice from your GP.

Treatment of PCS is usually about managing any ongoing symptoms. Specific treatment and therapies will be recommended depending on what the individual is experiencing.

It can be difficult to come to terms with everything following a brain injury and it's not always obvious what help you need. To make sure you don't suffer any further damage from PCS, it's important to seek help from people who truly understand the complexity of these types of injuries and the impact they can have.



# Diagnosing mild brain injuries

As we mentioned earlier, mild brain injuries can be very difficult to diagnose for a number of reasons. Many of the common symptoms associated with brain injuries can also be linked to other conditions, such as depression or chronic fatigue syndrome, and it's not uncommon for a person with a brain injury to be misdiagnosed.

Doctors aren't helped by scans, which don't pick up these injuries. Just because a brain injury may not have been properly recognised or diagnosed, it doesn't make the symptoms any less serious for the person who is injured.

After a brain injury, it can sometimes be hard to recognise any changes or explain your

symptoms fully. Family and friends may be able to help you, by discussing how your behaviour has changed.

It's a good idea to keep a diary of your symptoms and experiences to show to your GP. The charity Headway also has a useful [fact sheet](#) to help GPs understand your head injury and how it is affecting you. Requesting copies of your medical notes is also a good way to collect all your medical history and symptoms in one place. Asking for past medical notes can take some time and persistence, but will be worth it in the long run, help you to provide a more accurate medical history to any potential treating doctors.



# How a mild brain injury can impact your life

Mild brain injuries can have a big impact on many different aspects of your life - even if you don't recognise it at first. Here are some everyday things that may be affected by a head injury:



Driving



Alcohol



Sustaining another  
brain injury



Playing sports



Returning to  
work



Socialising

# How a mild brain injury can impact your life

## Driving

You should avoid driving until you feel completely well. Many brain injury symptoms, including issues with concentration, dizziness and balance problems, can all impact your ability to drive. When you do feel ready to drive, take things slowly and avoid long journeys. You might also need to inform the DVLA of any changes to your medical condition following a brain injury. Your GP will be able to advise you about this, or you can find out more on the [DVLA website](#).

## Alcohol

After a brain injury, people often notice their tolerance to alcohol is a lot lower. It is recommended you avoid alcohol immediately after a brain injury, and only consume alcohol again when you are feeling better. Drinking alcohol can also slow down brain injury recovery and make you more likely to sustain further injuries.

## Sustaining another brain injury

Knocks to the head have a cumulative effect, and sustaining other brain injuries can cause serious complications. You should avoid activities that increase the risk of injury to your head (such as playing contact sports) until you have fully recovered.



# How a mild brain injury can impact your life

## Playing sports

You should not play any contact sports for at least three weeks after a mild brain injury. If you return too soon, any further injuries to your head can be very dangerous. Many sports (such as football and rugby) have return to play protocols, but other sports are not as strict. You should always be vigilant about returning to play after a head injury. As Headway say – if in doubt, sit it out.

## Returning to work

Getting back to work after a brain injury can be a huge step. You might find certain tasks more challenging, or even feel unable to return to the job you did before. Speak to your employer about any worries you may have. They should be able to organise special arrangements to make your return to work as smooth as possible. You may want to consider a gradual return to work or to return on light duties. While you are recovering, try to avoid stressful situations as much as possible, and only return to work when you feel ready.



# How a mild brain injury can impact your life

## Socialising

The changes in your mood, as well as irritability, fatigue, and being overwhelmed by noisy environments can make socialising hard after a brain injury. Other people might not understand how your brain injury is impacting you or what difficulties you are facing. Even though it can be tricky, try not to shut yourself away. Explain your symptoms to loved ones and try to arrange social activities that are more suitable for you.

After a minor brain injury, these activities should become easier as you recover. If you're still struggling after a few weeks have passed, however, then your injury might be worse than first thought. If you suspect your injury is more serious than your doctor suggested, you should trust your instincts and seek advice from a medical professional with experience in mild TBIs



# Further support after a mild or moderate brain injury

Not having the right support after a brain injury can not only be frustrating - it can seriously impact the quality of your life. Without the help and attention you need, recovery can be extremely difficult.

While your GP may not be able to diagnose your brain injury, they can refer you to a specialist, such as a neurologist, who will be able to offer more specific assessment and treatment recommendations. However, waiting lists for these specialists can often be long.

You might want to visit a private neurologist or neuropsychologist to be assessed more quickly - a list of registered clinical neuropsychologists can be found on [The British Psychological Society](#) website.



# Charities and support groups

There are also some charities and support groups that can provide you with useful information, advice and guidance after a brain injury. These include:

## **Headway**

Headway is a national charity who provide support to brain injured people and their families. They also have a network of local groups and branches across the UK to provide support for people after a brain injury.

[www.headway.org.uk](http://www.headway.org.uk)

Tel: 0808 800 2244

## **BASIC**

The Brain and Spinal Injury Centre (BASIC) helps patients to access high-quality rehabilitation and treatment after a brain or spinal injury.

[www.basiccharity.org.uk](http://www.basiccharity.org.uk)

Tel: 0161 707 6441

## **Brain Injury Group (BIG)**

BIG provide support and information to people living with a brain injury, including injured people, their families and health and social care professionals.

[www.braininjuryisbig.org.uk](http://www.braininjuryisbig.org.uk)

Tel: 01483 770 999



# Helping friends and family to understand brain injuries

Sometimes, friends and family can find it hard to understand a brain injury and how this has impacted on your life. Brain injury is often referred to as an invisible injury, because the injured person may appear completely fine on the outside.

If you're struggling to explain your brain injury to friends and family, it's a good idea to ask them to read the information in this eBook. This can help them to understand mild brain injuries better, and the symptoms and problems you may be facing.

Try to explain to your loved ones how you are feeling and help them understand what they can do to assist with your recovery.



# About CFG Law

After a mild brain injury, there is a lot to consider, but the most important thing is having the support in place to make your road to recovery as quick and comfortable as possible.

At CFG Law, we understand how difficult and stressful it can be following a 'mild' TBI that may be more serious than first thought. Accessing the right treatment and understanding the impact this type of injury has on your day-to-day life is often confusing and difficult.

That's why we help our clients by offering more than expert legal advice. We work with medical professionals and charity organisations to help you get access to all the diagnosis, treatment options and funding you will need to get better. Our team of solicitors are not just experienced with traumatic brain injury cases, but dedicated to helping your recovery in any way they can.

Even if you think you do not need legal advice, we are happy to help you with any questions you may have and share our experiences with you.

**Get free advice  
from a specialist head  
injury solicitor**

